

17

DIVERSITY OF AROIDS (ARACEAE) IN PERLIS STATE PARK, PERLIS

Baharuddin Sulaiman & Mashhor Mansor

ABSTRACT

During the field collections, more than 20 aroid species were recorded and identified. They are *Aglaonema nitidum*, *A. simplex*, *Alocasia denudata* var. *denudata*, *A. lowii*; *Amorphophallus carneus*, *A. paeoniifolius*, *A. prainii*; *Arisaema roxburghii*, *A. wrayi*, *Colocasia esculenta* var. *esculenta*, *C. gigantea*; *Epipremnum giganteum*, *Homalomena coeruleascens*, *H. trapezifolia*, *Pothos scandens*, *P. latifolius*, *Raphidophora beccarii*, *R. foraminifera*, *Schismatoglottis calyptata* var. *calyptata*, *Scindapsus hederaceus*, *S. perakensis*, *Typhonium filiforme*, *T. flagelliforme*, *T. roxburghii*, and *T. trilobatum*. It should be noted that some of the species such as *Arisaema roxburghii*, *Arisaema wrayi*, *Raphidophora beccarii*, *Typhonium filiforme* and *Typhonium flagelliforme* are considered as rare.

ABSTRAK

Semasa pengutipan di lapangan lebih daripada 20 spesies keladi telah dirakam dan dicamkan. Meraka adalah *Aglaonema nitidum*, *A. simplex*, *Alocasia denudata* var. *denudata*, *A. lowii*; *Amorphophallus carneus*, *A. paeoniifolius*, *A. prainii*; *Arisaema*

roxburghii, *A. wrayi*, *Colocasia esculenta* var. *esculenta*, *C. gigantea*; *Epipremnum giganteum*, *Homalomena coeruleascens*, *H. trapezifolia*, *Pothos scandens*, *P. latifolius*, *Raphidophora beccarii*, *R. foraminifera*, *Schismatoglottis calyprata* var. *calyprata*, *Scindapsus hederaceus*, *S. perakensis*, *Typhonium filiforme*, *T. flagelliforme*, *T. roxburghii*, and *T. trilobatum*. Harus diingatkan bahawa kebanyakan spesies seperti *Arisaema roxburghii*, *Arisaema wrayi*, *Raphidophora beccarii*, *Typhonium filiforme* dan *Typhonium flagelliforme* adalah dianggap langka.

INTRODUCTION

Aroids are herbaceous monocotyledonous plants having heart-shape or cordiform, generally broad and pinnately veined leaves. The species are characterized by their inflorescences, which consist of a fleshy spadix surrounded by sessile small flowers and lack floral bracts. The inflorescence is covered by a specialized attractive organ called spathe.

The state of Perlis which is located in the northern part of Peninsular Malaysia and its prominent feature is a row of limestone hills with the formation of karst and razor sharp pinnacles. Setul and Chuping formations which are formed during geological era have created individual isolated hills and thus resulted in a characteristic distribution of vegetation especially the composition and diversity of the aroids. Generally the aroid species are found in a tropical areas and also are distributed world wide. This family is grouped into nine subfamilies, 106 genera and 3200 species (Croat, 1979, 1994). In Peninsular Malaysia, there are about 23 genera with 123 species (Ridley, 1925). The species are normally found in various habitats with special reference to wetlands; ranging from swamps, ponds, lakes, canals, rivers to rice fields. Some species thrive well in forest floors with good canopy coverage. To date, documents on limestone aroid are limited

A pioneer and comprehensive study on the Peninsular Malaysia Araceae had been conducted by Ridley (1925) and about 23 genera with 123 species. Other studies on the family are documented by Latiff *et al.* (1995), Ghani (1983, 1984), Neoh (1992) and Kress (1995). Mansor and Sulaiman (1997) have listed several species of aroid in swampy areas in Pondok Tanjung Forest Reserve. In addition, Sulaiman (1977) have surveyed riverine Araceae in Peninsular Malaysia.

METHODOLOGY

A field survey on the family Araceae was conducted at limestone hills in Perlis between 28 September 1999 to 5 October 1999. The plants were collected

and subsequently made into herbarium specimen for future references. The species were identified based on Henderson (1954), Ridley (1925), Bown (1988), Hay (1996a, 1996b), Hetterscheid (1996), Bogner and Nicolson (1991) and Mayo *et al.* (1997)

RESULTS AND DISCUSSION

A total of 14 Araceae genera were found in Perlis limestone hills and the genera are *Aglaonema*, *Alocasia*, *Amorphophallus*, *Amydrium*, *Anadendrum*, *Arisaema*, *Colocasia*, *Epipremnum*, *Homalomena*, *Pothos*, *Raphidophora*, *Schismatoglottis*, *Scindapsus* and *Typhonium*. Some of the genera such as *Aglaonema*, *Alocasia*, *Arisaema*, *Homalomena*, *Pothos*, *Scindapsus*, *Raphidophora*, *Schismatoglottis* and *Typhonium* have the ability to grow as terrestrial plants under the forest canopy. Only *Amorphophallus*, *Colocasia* and *Epipremnum* could adapt to higher light intensity which were the open places and disturbed areas. *Raphidophora*, *Scindapsus*, *Pothos* and *Epipremnum* are strictly climbers.

Based on the classification from Bogner and Nicolson (1991), the Araceae in limestone hills of Perlis can be grouped into five subfamilies namely Aroideae, Pothoideae, Philodendroideae, Monsteroideae and Colocasioideae (Table 1). The genera *Arisaema*, *Amorphophallus* and *Typhonium* are grouped in subfamily Aroideae. Two species of *Arisaema* are *A. roxburghii* and *A. wrayi*. *Amorphophallus carneus*, *Amorphophallus paeoniifolius* and *Amorphophallus prainii* were found in a more open sites. *Typhonium filiforme* could only survive growing in between the rocks on limestone hills.

Colocasioideae consist of genera *Colocasia* and *Alocasia*. Each genus is represented by two taxa in limestone hills namely *Colocasia esculenta* var. *esculenta* and *C. gigantea* and *Alocasia denudata* var. *denudata* and *A. lowii*. *Homalomena* and *Schismatoglottis* are grouped in subfamily *Philodendroideae*. Two species of *Homalomena* that are *H. coeruleascens* and *H. trapezifolia* have been recorded. Small colonies of *Schismatoglottis calyptata* var. *calyptata* are found in limestone hills.

Climbers are represented by *Amydrium humile*) and *Raphidophora beccarii* and *R. foraminifera* in the subfamily *Monsteroideae*. Only *Pothos scandens* and *P. latifolius* which are also climbers belong to the subfamily *Pothoideae*.

The Classification of Araceae based on inflorescences types is shown in Table 2. Eighteen species of Araceae found in limestone hills in Perlis have unisexual flower and 10 species have bisexual flower type. It should be noted that all unisexual flower species are ground flora whereas bisexual flowers are climbers.

Table 1: Classification of Araceae (After Bogner & Nicolson, 1991).

Subfamily	Tribe	Genera
Pothoideae		Pothos
Monsteroideae	Anadendreae	Anadendrum
	Monstereae	Amydrium Raphidophora Epipremnum Scindapsus
Philodendroideae	Philodendreae	Homalomena Schismatoglottis
	Aglaonemateae	Aglaonema
Colocasioideae	Colocasieae	Colocasia Alocasia
Aroideae	Thomsonieae	Amorphophallus
	Areae	Arisaema Typhonium

Table 2: Classification of Araceae based on inflorescence and flower types.

Unisexual flower

Arisaema roxburghii and *A. wrayi*

Amorphophallus prainii, *A. carneus* and *A. paeoniifolius*

Alocasia denudata var. *denudata* and *A. lowii*

Aglaonema nitidum and *A. simplex*

Colocasia esculenta var. *esculenta* and *C. gigantea*

Homalomena coeruleascens and *trapezifolia*

Schismatoglottis calyptata var. *calyprata*

Typhonium filiforme, *T. roxburghii*, *T. flagelliforme* and *T. trilobatum*

Bisexual flowers

Amydrium humile

Anadendrum montanum

Epipremnum giganteum

Pothos scandens and *P. latifolius*

Raphidophora beccarii and *R. foraminifera*

Scindapsus perakensis, *S. hederaceus* and *S. perakensis*

CONCLUSION

More intensive study is needed especially on the ecological aspect and also the adaptation of the family to the limestone areas.

ACKNOWLEDGEMENTS

The authors are grateful to Universiti Sains Malaysia, Penang for providing a Research Grant No. 304/Pbiologi/ 633094.

REFERENCES

- Bogner, J. & D. H. Nicolson. (1991) A revised classification of Araceae with dichotomous keys. *Willdenowia* 21: 35-50.
- Bown, D. (1988) *Aroids, plants of the Arum family*. Century Publication. 256 pp.
- Croat, T. B. (1979) The distribution of Araceae. In: K. Larsen & L. B. Holm-Nielsen (eds.), *Tropical botany*. Academic Press, London. Pp. 291-308.
- Croat, T. B. (1994) Taxonomic status of Neotropical Araceae. *Aroideana* 17: 33-60.
- Ghani, F. D. (1983) Ornamental and edible aroids of Peninsular Malaysia. *Aroideana* 6: 129-132.
- Ghani, F. D. (1984) The potentials of aroids in Malaysia. In: S. Chandra (ed.), *Edible aroids*. Clarendon Press, Oxford. Pp. 10-16.
- Hay, A. (1996a) The genus *Schismatoglottis* Zoll. & Moritz (Araceae: Schismatoglottideae) in Peninsular Malaysia and Singapore. *Sandakania* 7: 1-30.
- Hay, A. (1996b) A new Borneon species of *Colocasia* Schott (Araceae: Colocasieae), with a synopsis of the genus in Malesia and Australia. *Sandakania* 7: 31-48.
- Henderson, M.R. (1954) Malayan wild flowers. *Monocotyledons*. Malayan Nature Society, Kuala Lumpur. Pp. 213-243.
- Hetterscheid, W. & S. Ittenbach. (1996) Everything you always wanted to know about *Amorphophallus*, but were afraid to stick your nose into! *Aroideana* 19: 7-131.
- Kress, M. (1995) *Medicinal chest from Malaysian jungles*. Wings of Gold, Kuala Lumpur. Pp. 16-22.
- Latiff, A., K. Mat Salleh, A. Zainudin Ibrahim, S. Muzni, I.M. Turner & J.W. Yong. (1995) Preliminary checklist of flowering plants from Temenggor Forest Reserve, Hulu Perak, Malaysia. *Mal. Nat. J.* 48: 175-188.
- Mansor, M. & B. Sulaiman. (1997) Notes on Araceae of the Pondok Tanjung Forest Reserve, Peninsular Malaysia. *Tropical Biodiversity* 4: 157-161.
- Mayo, S. J., Bogner, J. and Boyce, P. C. (1997). The Genera of Araceae. Royal Botanic Gardens. Kew. 370pp.

- Neoh, C. K. (1992) *Typhonium divaricatum* (Rodent tuber): A promising local plant in the fight against cancer. Med. J. Mal. 47: 86-89.
- Ridley, H. N. (1925) *The Flora of Malay Peninsula*. Vol. V. L. Reeve & Co. Ltd., London. 470 pp.
- Sulaiman, B. (1997) *Taksonomi dan diversiti Araceae di sungai-sungai Semenanjung Malaysia*. MSc. Thesis, University of Science Malaysia. 105 pp.